

Journal of Depression and Anxiety Forecast

The Combined Use of Carbamazepine and Lithium in the Treatment of Outpatients with Rapid Cycle Bipolar Disorder without Psychiatric Comorbidity: Randomized, Double-Blind Placebo-Controlled Study

Aliyev NA^{1*} and Aliyev ZN²

¹Department of Psychiatry and Drug Addiction, Azerbaijan State Advanced Training Institute, Baku, Azerbaijan

²Department of Psychiatry, Azerbaijan Medical University, Baku, Azerbaijan

Abstract

Objective: Rapid cycle bipolar disorder may be more resistant to treatment than other bipolar types. However, the effect of the combined use of carbamazepine and lithium in the treatment of outpatients with rapid cycle bipolar disorder without psychiatric comorbidity has not been studied in a double-blind placebo-controlled design.

Materials and Methods: Eligible participants to meeting the DSM-5 criteria for rapid cycle bipolar disorder was a double-blind, placebo-controlled trial, were required to be between 18 and 65 years. Response and side effects with carbamazepine and lithium and placebo were compared by using analysis of variance (ANOVA) and chi-square tests. The responder was conducted by X² demonstrated superior for carbamazepine and lithium than for placebo. Carbamazepine and lithium was generally well tolerated by the patients in the study, although more carbamazepine and lithium-treated patients (n=2) discontinued the study early because of adverse events.

Results: Twenty seven of the 36 combined use of carbamazepine and lithium-treated participants responded by 12 weeks, *versus* six of the 38 placebo-treated participants! ($p < 0.001$). The two common side effects leading to discontinuation in the carbamazepine and lithium group were allergic reaction and drowsiness. Sweating and headache frequent complaint during placebo treatment occurred in two of 38 men. There were not unexpected or serious adverse events.

Conclusions: The combined use of carbamazepine and lithium in the treatment of outpatients with Rapid cycle bipolar disorder demonstrated superior for carbamazepine and lithium than for placebo.

Keywords: Rapid cycle bipolar disorder; Carbamazepine; Lithium; Treatment

Introduction

According to the epidemiological data the prevalence of bipolar disorder DSM-5 over the course of 12 months and throughout life was 1.5% and 2.1% and did not differ between men (1.6% and 2.2%) and women (1.5% and 2, 0%) [1].

Total spending on BDI in 2015 was estimated at \$202.1 billion, which corresponds to an average of \$81,559 per person, while excess spending on BDI was estimated at \$ 119.8 billion, which corresponds to an average of 48333 USD per person [2].

The total cost of mental illness is estimated at more than 4% of GDP-or more than 600 billion Euros in 28 EU countries. Severe mental illness, such as bipolar disorder, affects nearly 5 million people (1.0% of the population), while schizophrenic disorders affect another 1.5 million people (0.3%) [3].

Available literature indicates that the pharmacological treatment of fast cycling is still scarce, and therefore there is no consensus on its optimal pharmacological treatment. Clinical trials specifically studying rapid cycling are needed to unravel the proper management of bipolar disorder with rapid cycling [4]. Some authors conclude that rapid cycling is a common, albeit insufficiently recognized, condition for bipolar disorder, and it represents a worsening of the primary disorder.

OPEN ACCESS

*Correspondence:

Nadir A. Aliyev, Department of Psychiatry and Drug Addiction, Azerbaijan State Advanced Training Institute, Baku, Azerbaijan.

E-mail: aliyevnadir@yahoo.com

Received Date: 29 Apr 2020

Accepted Date: 15 May 2020

Published Date: 21 May 2020

Citation: Aliyev NA, Aliyev ZN. The Combined Use of Carbamazepine and Lithium in the Treatment of Outpatients with Rapid Cycle Bipolar Disorder without Psychiatric Comorbidity: Randomized, Double-Blind Placebo-Controlled Study. *J Depress Anxiety Forecast.* 2020; 3(1): 1013.

ISSN 2643-7139

Copyright © 2020 Aliyev NA. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

There is no convincing evidence that fast cycling is a separate subtype. Early recognition of this pattern can lead to an improvement in treatment strategy and an improvement in the long-term course. Conceptualizing fast cycling according to the results of these studies should be an important step forward [5].

Pharmacotherapy for rapid cycle bipolar disorder is an active area of research. A variety of drug groups have been shown to be effective in treating many of the rapid cycle bipolar disorder. From a therapeutic point of view, a previous study showed the effectiveness of several drugs. For example, lamotrigine is known to increase euthymia in patients with RCBD compared with placebo [6]. In addition, pramipexol has been shown to facilitate resistant depression in patients with RBC [7]. In patients with concomitant pathological conditions, such as hypothyroidism, levothyroxine supplementation has been shown to eliminate the symptoms of RCBD in patients [8]. Finally, it has also been shown that ketamine is extremely useful in patients with RBC with active suicidal ideation [9]. Bipolar disorder is characterized by episodes of mood that almost always recur. Patients who experience at least four episodes over a 12 month period are classified as “rapid cycles”. Patients who experience at least four episodes during a 12 month period are classified as “rapid cycling” [10]. The term was first used to describe bipolar patients who did not respond to lithium [3]. However, it is now clear that all drugs (including lithium) are less effective for patients with “rapid cycles” than patients with non-“rapid cycles” [9].

Rapid cycle bipolar disorder may be more resistant to treatment than other bipolar types. In the open phase of the two-phase study, 133 depressed patients with rapid cycle bipolar disorder began to take lithium plus divalproex. Only 14% stabilized; 10% withdrew due to side effects; 17% were not committed. In the double-blind phase, 49 respondents were not randomized to placebo or lamotrigine. The two groups showed no differences in response rates [3].

Studies show that the rapid cycle is different from other forms of bipolar disorder. People with such mood changes may respond differently to standard and experimental treatments than other people with bipolar disorder. With his sudden and unpredictable mood changes, rapid cycles can be more difficult than other types of bipolar disorder.

For rapid cycling (can be used for bipolar disorder I or bipolar disorder II): at least four episodes of mood during the previous 12 months that meet the criteria for a manic, hypomanic or major depressive episode. Episodes are limited to partial or complete remission for at least 2 months or a transition to an episode of opposite polarity (for example, an episode with major depression to a manic episode). An essential feature of bipolar disorder with rapid cycling is the occurrence of at least four episodes of mood over the previous 12 months. These episodes can occur in any combination and order. Episodes should meet the criteria for both the duration and the number of symptoms for a major depressive, manic, or hypomanic episode and should be limited to either a period of complete remission or a transition to an episode of opposite polarity. Manic and hypomaniac episodes are considered to be at the same pole. Except for the fact that they occur more often, the episodes that occur in the fast-cycle pattern are no different from the episodes that occur in the non-rapid-cycle mode. The episodes of mood that are taken into account when determining the fast cycle pattern exclude those episodes that are directly caused by the substance (for example, cocaine, corticosteroids) or another disease [10].

The crucial issue is demonstrated by reviewing the literature that the effect of the combined use of carbamazepine and lithium in the treatment of outpatients with Rapid cycle bipolar disorder without psychiatric comorbidity has not been studied in a double-blind placebo-controlled design. This study reports on a randomized, double-blind, placebo-controlled of carbamazepine and lithium in the treatment of outpatients with Rapid cycle bipolar disorder. The aim of this work was to study the effectiveness of carbamazepine in combination with lithium in the treatment of rapid- cycle bipolar disorder in the framework of type I bipolar disorder.

Materials and Methods

This was a double-blind, placebo-controlled trial for patients diagnosed with DSM-5 for rapid cycle bipolar disorder. The patients gave their informed, written consent to participate.

In accordance with the Helsinki Declaration of the World Medical Association “Recommendations for doctors engaged in biomedical research involving people”, adopted by the 18th World Medical Assembly (Finland, 1964, revised in Japan in 1975, Italy-1983, Hong Kong-1989, the South African Republic-1996, Edinburgh-2000); The Constitution of the Republic of Azerbaijan, the Law “On Psychiatric Assistance” (adopted on 12.06.2001, with amendments and additions -11.11.2011, Decisions of the Cabinet of Ministers of the Republic of Azerbaijan No. 83, dated April 30, 2010 “On Approval of the Rules for Conducting Scientific, Preclinical and Clinical studies of medicines” are established. The conditions of the conducted researches corresponded to the generally accepted norms of morality, the requirements of ethical and legal norms, as well as the rights, interests and personal dignity of the participants of the studies were observed.

- a) Conducted research is adequate to the topic of research work.
- b) There is no risk for the subject of research.
- c) Participants in the study were informed about the goals, methods, expected benefits of the study and associated with risk and inconvenience in the study.
- d) The subject’s informed consent about participation in the research was received.

The decision of the Ethical Committee at the Azerbaijan Psychiatric Association on the article of NA. Aliyev, Z.N. Aliyev “A Double-Blind Placebo-Controlled Trial of The combined use of carbamazepine and lithium in the treatment of outpatients with Rapid cycle bipolar disorder without psychiatric comorbidity: Randomized, double-blind placebo-controlled study” submitted for publication in psychiatric journals: in connection with compliance with its legislative requirements and regulatory documents is to approve the article by N.A. Aliyev, Z.N. Aliyev “The combined use of carbamazepine and lithium in the treatment of outpatients with Rapid cycle bipolar disorder without psychiatric comorbidity: Randomized, double-blind placebo-controlled study”. Patients were observed at the Mental Health Center of the Ministry of Health of the Republic of Azerbaijan. The study was conducted from January 1, 2017 to January 2018.

Also we excluded sexually active subjects with active or unstable epilepsy, other genetic syndromes or congenital infections associated with autistic-like syndromes, prematurity; subjects who have been treated within the previous 30 days by any medication known to have a clearly defined potential for toxicity or with any psychotropic drugs;

subjects with clinically significant abnormalities in laboratory tests or physical examination; subjects with a history of hypersensitivity or serious side effects associated with the use of carbamazepine and lithium, and subjects who, during the previous 3 months, started new non-pharmacological procedures, such as diet, vitamins and psychosocial therapy. A detailed clinical interview with parents by a clinical expert, accompanied by physical examination and blood analysis, was used to ensure that subjects did not meet any exclusion criteria.

This was a double-blind, placebo-controlled trial for patients diagnosed with DSM-5 for rapid cycle bipolar disorder. A structured clinical interview, for DSM-5 Axis I Disorder, Patient Edition, was used to diagnose rapid cycle bipolar disorder according to DSM-5. Eighty patients (40 men and 40 women) whom we studied were under observation in Mental Clinic for Outpatients of Baku City of Azerbaijan Republic. The length of the washout was 2 weeks. Patients were washout from the all medications. The method of randomization was given by lottery. Each patient was randomized to receive either combined use of carbamazepine and lithium (40 patients) accordingly, in doses of 200 mg three times and 300 mg three times per day for 12 weeks or matched placebo (48 patients) in a double-blind manner. All patients were evaluated by experienced clinicians. The average level of concentration of carbamazepine and lithium in the blood plasma in patients was $7.8 \pm 5.9 \mu\text{g/ml}$ and $0.6\text{--}0.8 \text{ mmol/l}$, respectively, range.

Eligible participants, in addition to meeting the DSM-5 criteria for rapid cycling bipolar disorder and were required to be between 18 and 65 years of age. We excluded serious medical conditions including with other psychiatric disorders (e.g. bipolar disorder II type, schizophrenia, patients judged to be at serious suicidal or homicidal risk, dependence of psychoactive drugs, somatic, neurological illness etc). Also we excluded unstable epilepsy, other genetic syndromes or congenital infections associated with autistic-like syndromes, prematurity; subjects who have been treated within the previous 30 days by any medication known to have a clearly defined potential for toxicity or with any psychotropic drugs; subjects with clinically significant abnormalities in laboratory tests or physical examination; subjects with a history of hypersensitivity or serious side effects associated with the use of carbamazepine and lithium, and subjects who, during the previous 3 months, started new non-pharmacological procedures, such as diet, vitamins and psychosocial therapy. A detailed clinical interview with parents by a clinical expert, accompanied by physical examination and blood analysis, was used to ensure that subjects did not meet any exclusion criteria.

Patients clinically significant of abnormal laboratory or EEG findings were ineligible. Patients before the study had used antidepressants, antipsychotics, anxiolytics, benzodiazepines, SSRI and venlafaxine. Washout of all medicines was two weeks.

Analysis of response refers to the last observation carried forward for all subjects who had valuable efficacy at baseline and with treatment. The responder analysis was conducted by using the chi-square (χ^2) and analysis of variance (ANOVA) according to Glantz [11].

Results

Characteristics of the patients randomly assigned to the two treatments are shown in Table 1.

Statistical differences between the two groups are not significantly.

Table 1: Sociodemographic characteristics of patients with rapid cycle bipolar disorder.

Characteristic	Carbamazepine + lithium n =40	Placebo n = 40
	Mean SD	Mean SD
Age (years)	38.0 ± 10.2	37.4 ± 9.8
Duration of illness	10.2 ± 6.0 n = 40%	10.0 ± 5.8 n = 40%
Education:		
Primary school	28 (70)	30 (75)
Secondary school	12 (30)	10 (25)
Marital status:		
Never married	10 (25)	10 (25)
Married	20 (50)	18 (45)
Divorced or separated	10 (25)	12 (30)
Employment status:		
Unemployed	18 (45)	20 (50)
Employed	22 (55)	20 (50)

Note: Differences between groups are not significant.

Table 2: The results of the square analysis.

Treatment Groups	Observed number	Expected number	chi-square (χ^2)
	Yes improvement	Not improvement	Total
Placebo	6 (16.43)	32 (21.57)	38
Carbamazepine + lithium	27 (15.56)	9 (20.43)	36
Total	33	41	74

Note: Expected numbers indication in the brackets. $\chi^2=22.68$, $df=1$, $p<0.001$.

The results of treatment are shown in Table 2. As shown from the Table 2, six patients did not return for at least one subsequent assessment, leaving 74 patients (36 taking carbamazepine and lithium and 38 taking placebo) in the valuable study group.

The responder was conducted by χ^2 demonstrated superior for carbamazepine and lithium than for placebo (Table 2). Carbamazepine and lithium was generally well tolerated by the patients in the study, although more carbamazepine and lithium-treated patients ($n=2$) discontinued the study early because of adverse events. The two common side effects leading to discontinuation in the carbamazepine and lithium group were allergic reaction and drowsiness. Sweating a frequent complaint during placebo treatment occurred in two of 38 men. There were not unexpected or serious adverse events.

Discussion

History and Discovery, structure-activity relationships, pharmacological profiles, mechanisms of action, pharmacokinetics and disposition, indications and efficacy response, side effects and toxicology, drug-drug interactions lithium and carbamazepine are described in detail in the literature [9]. Therefore, we will not dwell on them.

To our knowledge, this is the first report of a randomized, double-blind, placebo-controlled study of a Carbamazepine+lithium in the treatment of outpatients with Rapid cycle bipolar disorder without psychiatric comorbidity. Our data suggest that Carbamazepine+lithium is efficacious in the management of rapid cycling bipolar disorder, as the participants had a clinically and statistically significant improvement in rapid cycling bipolar disorder over 12 weeks of treatment.

From the eighty patients six patients did not return for at least one subsequent assessment, leaving 74 patients (36 Carbamazepine+lithium and 38 taking placebo) in the valuable study group. Twenty six (94%) of the 36 Carbamazepine+lithium treated participants responded by 12 weeks, versus six (16%) of the 38 placebo-treated participants. The most common and problematic side effect in the carbamazepine and lithium group was on one allergic reaction and drowsiness. As a side effects in the placebo group indicated headache and sleeplessness.

As noted at the beginning pharmacotherapy for rapid cycle bipolar disorder is an active area of research. Pharmacotherapy for rapid cycle bipolar disorder is still very relevant [12-14].

Our data partially coincide with some results of studies by other authors [15,16]. It was found that carbamazepine has very limited possibilities in the treatment of rapid cycles, especially in the form of monotherapy. It has moderately pronounced effectiveness in the manic phase and weakly moderately effective in the depressive phase [17].

At least 8 placebo-controlled randomized trials have demonstrated that lithium is a prophylactic in bipolar disorder [13]. A meta-analysis of 16 studies of bipolar disorder conducted by Tondo and colleagues [9]. Tondo L, Hennen J, Baldessarini RJ showed that lithium can be considered first-line treatment in the prevention of rapid cyclic bipolar disorder. Many studies confirm the effectiveness of lithium as a mood stabilizer [4,9,18-21].

Despite the obvious limitation of this study due to the small number of samples, we can argue that it shows promising data on the usefulness of the combined use of carbamazepine and lithium in the treatment of outpatients with Rapid cycle bipolar disorder in the routine treatment of complex pathologies in this illness. It will be important to explore further effects of the combined use of carbamazepine and lithium in the treatment of rapid cycle bipolar disorder.

Conclusion

As indicated earlier, rapidly recurring bipolar disorders are difficult to treat. Treatment should be based on current practical guidelines and expert advisory groups on the treatment of bipolar disorder. Historically, rapid cycles usually did not respond to lithium or carbamazepine. The long-term result of lithium prophylaxis in bipolar disorder: a 5-year prospective study of 402 patients in a lithium clinic. Data from the largest prospective, placebo-controlled, rapid-cycle bipolar disorder study demonstrated the efficacy of lamotrigine in patients with euthymia or mood episodes. Used a 6-week titration of lamotrigine to 200 mg/day. The results of these data showed that lamotrigine monotherapy was a useful treatment, especially for patients with rapid-cyclic bipolar II disorder [3,13]. More recent comparisons of lithium and valproic acid show no significant differences in efficacy between lithium and divalproate for long-term treatment of rapid-cycle bipolar disorder, but have shown that lamotrigine has the potential to complement both drugs in the treatment of depression [22].

Our Follow-up observation (catamnesus) was conducted from January 01, 2018 to July 01, 2019. The results of follow-up observations showed that patients receiving carbamazepine in combination with lithium are in a state in full remission during for 2.5 years.

Recommendation

1. As indicated earlier, rapidly recurring bipolar disorders are difficult to treat. Treatment should be based on current practical guidelines and expert advisory groups on the treatment of bipolar disorder.
2. The combined use of carbamazepine and lithium in the treatment of outpatients with Rapid cycle bipolar disorder demonstrated superior for carbamazepine and lithium than for placebo.
3. Carbamazepine and lithium - drug interactions must be considered.
4. Availability a history of allergic reactions must be considered.

Acknowledgment

The authors would like to thank staff of the Mental Health Center of the Ministry of Health of the Republic of Azerbaijan.

References

1. Blanco C, Compton WM, Saha TD, Goldstein BI, Ruan WJ, Huang B, et al. Epidemiology of DSM-5 bipolar I disorder: Results from the National Epidemiologic Survey on Alcohol and Related Conditions-III. *J Psychiatr Res.* 2017; 84: 310-317.
2. Cloutier M, Greene M, Guerin A, Touya M, Wu E. The economic burden of bipolar I disorder in the United States in 2015. *J Affect Disord.* 2018; 226: 45-51.
3. Maj M, Pirozzi R, Magliano L, Bartoli L. The long-term result of lithium prophylaxis in bipolar disorder: a 5-year prospective study of 402 patients in a lithium clinic. *Am J Psychiatry.* 1998; 155: 30-35.
4. Fountoulakis KN, Kontis D, Gonda X, Yatham LN. A systematic review of the evidence on the treatment of rapid cycling bipolar disorder. *Bipolar Disord.* 2013; 15: 115-137.
5. Carvalho AF, Dimellis D, Gonda X, Vieta E, McIntyre RS, Fountoulakis KN. Rapid cycling in bipolar disorder: a systematic review. *J Clin Psychiatry.* 2014.
6. Goldberg JF, Bowden CL, Calabrese JR, Ketter TA, Dann RS, Frye MA, et al. Six-month prospective life charting of mood symptoms with lamotrigine monotherapy versus placebo in rapid cycling bipolar disorder. *Biol Psychiatry.* 2008; 63: 125-130.
7. Aditya H, Aditi S, Malvika R, Janardhanan CN, Suresh BM. Pramipexole in the treatment of refractory depression in a patient with rapid cycling bipolar disorder. *Indian J Psychol Med.* 2015; 37: 473-474.
8. Chen P, Huang Y. Remission of classic rapid cycling bipolar disorder with levothyroxine augmentation therapy in a male patient having clinical hypothyroidism. *Neuropsychiatr Dis Treat.* 2015; 11: 339-342.
9. Viguera AC, Baldessarini RJ, Tondo L. Response to lithium maintenance treatment in bipolar disorders: comparison of women and men. *Bipolar Disord.* 2001; 3: 245-252.
10. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5). American Psychiatric Association. 2013.
11. Glantz SA. *Primer of Biostatistics.* 2012.
12. Huber J, Burke D. ECT and lithium in old age depression-Cause or treatment of rapid cycling? *Australas Psychiatry.* 2015; 23: 500-502.
13. Maj M, Pirozzi R, Formicola AMR, Tortrella A. Reliability and validity of four alternative definitions of bipolar disorder with fast cycling. *Am J Psychiatry.* 1999; 158: 303-305.

14. Sampath H, Sharma I, Dutta S. Treatment of suicidal depression with ketamine in rapid cycling bipolar disorder. *Asia Pac Psychiatry*. 2016; 8: 98-101.
15. Goldberg JF, Bowden CL, Calabrese JR, Ketter TA, Dann RS, Frye MA, et al. Six-month prospective life charting of mood symptoms with lamotrigine monotherapy versus placebo in rapid cycling bipolar disorder. *Biol Psychiatry*. 2008; 63: 125-130.
16. Jonathan S, Kemp DE. Treatment Choices for Resistant Bipolar Disorder. *Bipolar Disord*. 2012.
17. Calabrese JR, Fatemi SH, Kujawa M, Woyshville MJ. Predictors of response to mood stabilizers. *J Clin Psychopharmacol*. 1996; 16: 24-31.
18. Baldessarini RJ, Tondo L, Hennen J, Viguera AC. Is lithium still worth using? An update of selected recent research. *Harv Rev Psychiatry*. 2002; 10: 59-75.
19. Calabrese JR, Shelton , Rapport DJ, Youngstrom EA, Jackson K, Bilali S, et al. A 20-month, double-blind, maintenance trial of lithium *versus* divalproex in rapid-cycling bipolar disorder. *Am J Psychiatry*. 2005; 162: 2152-2161.
20. Koukopoulos A, Reginaldi D, Laddomada P, Floris G, Serra g, Tondo L. Course of the manic-depressive cycle and changes caused by treatments. *Pharmacopsychiatry*. 1980; 13: 156-167.
21. Health at a Glance: Europe 2018. State of Health in the EU Cycle. OECD Publishing. 2018.
22. Calabrese JR, Rapport DJ, Youngstrom EA, Jackson K, Bilali S, Findling RL. New evidence on the use of lithium, divalproate and lamotrigine in fast cycles of bipolar disorder. *Eur Psychiatry*. 2005; 20: 92-95.