

# Journal of Gynecology and Obstetrics Forecast

## Uvula Oedema after Application of Nitrous Oxide during Labour

Šanjug J<sup>1\*</sup>, Hrgović Z<sup>2</sup>, Jaška S<sup>3</sup>, Fureš R<sup>1-4</sup> and Gradicač M<sup>5</sup>

<sup>1</sup>Zabok General Hospital and Croatian Veterans Hospital, Zabok, Croatia

<sup>2</sup>Faculty of Dental Medicine and Health Osijek, Josip Juraj Strossmayer University of Osijek, Croatia

<sup>3</sup>Faculty of Medicine Osijek, Josip Juraj Strossmayer University in Osijek, Croatia

<sup>4</sup>Croatian Society for Pelvipereology, Croatia

<sup>5</sup>Johann Wolfgang Goethe- University, Frankfurt am Main, Germany

### Letter to the Editor

We would like to share our experience of uvula oedema during labour after application of inhaled nitrous oxide.

A 31 year old gravida 2 para women with neat family and obstetric history. During second pregnancy, gestational diabetes mellitus was diagnosed in 24<sup>th</sup> gestational week, regulated by diet [1].

Patient was hospitalized at 39<sup>th</sup> gestational week due to planned labour. Labour induction started at 1, 5cm dilated cervixes by PgE2 vaginal gel application. Six hours later amnion rupture happens spontaneously and when cervices were dilated 5cm analgesia started with inhaled nitrous oxide and infusion of oxytocin (5i.u. /ml).

Two hours later symptoms occurred as difficulty in speaking and swallowing, choking feeling and urge to vomiting. A doubled enlarged, oedematous, reddish uvula was observed without tongue oedema and enlarged tonsils.

That time cervices were completely opened and contraction frequency was 2/10 min with normal CTG values.

Treatment of uvula oedema started with oxygen application and corticosteroid therapy. Blood pressure (RR 130/80 mmHg), pulse 100bpm and oxygen saturation (98%) were monitored. Anaesthesiologist was also consulted. Examination of the respiratory system was unremarkable [2].

Fifteen minutes after the appearance of the first symptoms of uvula oedema started labour which finished successfully and male infant was born (4.170g, 52cm) with Apgar score 9/10 and umbilical arterial pH 7.31.

Response to the therapy started half hour later. After six hours symptoms of difficulties in speaking, swallowing and feeling of choking were minimized but uvula edema was still present but decreased by 1/3. Patient was monitored in ICU 24h after delivery and uvula oedema has with drowned within 24 hours. Patient left the hospital three days after delivery.

Complete blood count, coagulation, routine biochemical, liver function tests and CRP were done half hour after delivery. Deviations from normal values shown CRP=53.3mg/l and L=30.6x10<sup>9</sup>/L and which were normalized within 24 hours.

Uvula oedema symptoms were treated with methyl-prednisolon 125mg i.v, chlorpiraminchlorid 20mg, adrenalin 1:1000, calcium-gluconate 10ml [3].

Acute uvula oedema is rare occurrence and could be potentially life-threatening condition involving go the rupper airway structures such as apiglottis, causing rapid partial or complete airway obstruction and difficulties in speaking and breathing as it affects the vocalcord. Early and accurate diagnose and treatment is exceptionally important.

Etiology of isolated angio-oedema is usually due to type 1 hypersensitivity reaction and angio-oedema of uvula is not associated with other hypersensitivity reaction such as pruritis, skinrash, Hypotension or tachycardia. It is localised, recurrent, non-pruritic subcutaneous swelling that

### OPEN ACCESS

#### \*Correspondence:

Jadranka Šanjug, Zabok General Hospital and Croatian Veterans Hospital, Bračak 8, 49210 Zabok, Croatia.

**E-mail:** jadranka.sanjug@gmail.com

**Received Date:** 05 Apr 2019

**Accepted Date:** 23 Apr 2019

**Published Date:** 29 Apr 2019

**Citation:** Šanjug J, Hrgović Z, Jaška S, Fureš R, Gradicač M. Uvula Oedema after Application of Nitrous Oxide during Labour. *J Gynecol Obstet Forecast*. 2019; 2(1): 1009.

**Copyright** © 2019 Šanjug J. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

usually appears rapidly and resolves spontaneously.

The possible causes of recurrent uvula oedema are: idiopathic, hereditary, trauma, infections, allergy, medications and obstructive sleep apnea [4].

In our case, there is possibility of isolated uvula oedema occurrence due to nitrous oxide use in labour analgesia.

There is a possibility of oxidation of nitrous oxide to nitric oxide by hydrochloric acid due to urge to vomiting and aspiration of nitrous oxide. Nitric acid is a potent vasodilator and might cause uvula angio-oedema.

Oedema is thought to result of inflammatory induced increases in airway micro vascular permeability. Inflammatory mediators released in the airway mucosa include histamin, leucotriens and PAF are potent inducers of increased oedema and airway mucosal swelling.

It is unclear whether our patient's uvula oedema was result of an antigen response to nitrous oxide or is related to trauma caused

by dehydration due to exposure to dry gas and improper breathing technique (too deep or open mouth breathing).

## References

1. Alcoceba E, Gonzalez M, Gaig P, Figuerola E, Auguet T, Olona M. Edema of the Uvula: Etiology, Risk Factors, Diagnosis, and Treatment. *J Investig Allergol Clin Immunol.* 2010; 20: 80-83.
2. Evans TC, Roberge RJ. Quincke's disease of the uvula. *Am J Eerg Med.* 1987; 5: 211-216.
3. Partel RV, Cho C, Medd C, Cresswell J. Isolated non-hereditary angioneurotic oedema of uvula (Quincke's disease) in an adolescent. *BMJ Case Rep.* 2014.
4. Ford R, Bird S. Uvula oedema after prolonged use of inhald nitrous oxide during labour in an obstetric patient. 2015.