

SF Journal of Neurological Disorders and Rehabilitation

Graphomotor Skills in Children with Language Deficits in Primary School Age - Formation and Development

Spassova K*

Department of Public Health, Medical University of Varna, Varna, Bulgaria

Abstract

Introduction: The formation and the development of graphomotor skills in children with language deficits in primary school age is a difficult and long process, and its disturbance leads to serious problems not only with the process of literacy but also disrupt the purely academical learning.

Purpose: The aim of our study is to show the level of formation of graphomotor skills in children with language deficits in primary school age before and after the implementation of a system of therapeutic techniques, approaches and methods based on both the world's and Bulgarian's experience in the area of therapy of disturbance in graphomotor skills.

Method: The study encompasses a total of n=40 children in primary school, aged 7-8, who were subjected to logopedic therapy aimed to overcome difficulties in graphomotor skills. It was conducted for a period of 5 years. The therapy was done either individually or in groups of 2 children.

Results: The data from test 1 and test 2 show development and maturation of structures in the left hemisphere: frontoparietal and basal temporal, responsible for functioning of the spatial orientation and ideomotor praxis. For this reason the 7-8 years age group can be considered sensible for development of graphomotor skills as a component of formation of the written form of language and is especially suitable for intensive logopedic therapy aimed for its formation.

Conclusion: The implemented therapeutic techniques during the logopedic therapy are aimed not only for establishing the occulo-spacial orientation but also for formation and development of language competence, which are directly linked to formation of the written form of language.

Keywords: Graphomotor skills; Frontoparietal structures; Ideomotor praxis

Introduction

The formation and development of graphomotor skills in children with language deficits in primary school age is a hard and a long process and its disturbance leads to serious problems not only with the process of literacy but also disrupt the purely academical learning. In this particular age the establishment of literacy, as one of the important processes in human life, starts. This process is dependent on the normal development of senses (vision, hearing and touch), intellect and social environment.

The scientific work in this area show that the formation of the written skills is disrupted by disturbance in the visual- motor coordination in children and school pupils as a result of immaturity or false formation of some of the higher cortical functions which are responsible for development of the graphomotor skills and habits. Their disturbed formation leads to aberration in orientation in space and time, low motivation and insufficient self-control, expressed in difficulty in the process of formation of graphomotor skills and mastering the written form of language [1-3].

From psycho-physiological view and based on neuro-psychological model in the process of mastering written language three features take part: visual, kinetic and auditory. Historically speaking the leading role has the discovered by Exner "area of writing" in the posterior part of the second frontal gyrus, the so called premotor field or Brodmann area 8 [1,4].

Purpose of our study is to show the level of formation of graphomotor skills in children with language deficits in primary school age (7-8 years) before and after the implementation of a system of therapeutic techniques, approaches and methods based on both the world's and Bulgarian's experience in the area of therapy of disturbance in graphomotor skills.

OPEN ACCESS

*Correspondence:

Kalinka Spassova, Department of Public Health, Medical University of Varna, Varna, Bulgaria.

E-mail: spasova.kalina@abv.bg

Received Date: 30 Apr 2020

Accepted Date: 22 May 2020

Published Date: 28 May 2020

Citation: Spassova K. Graphomotor Skills in Children with Language Deficits in Primary School Age - Formation and Development. *SF J Neurol Disord Rehabil.* 2020; 1(1): 1002.

Copyright © 2020 Spassova K. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Table 1:

Group	2012/2013	2013/2014	2014/2015	2015/2016	2016/2017	Total
7-8 y.	n=8	n=10	n=7	n=9	n=6	n=40

Table 2:

Methods	Mean	Total	Standard deviation	Standard error of the mean
Head 1entry	7.1375	40	0.86963	0.13750
Head 1exit	7.5250	40	0.68827	0.10883
Head 1.2. entry	5.9375	40	0.64239	0.10157
Head 1.2. exit	6.5125	40	0.59364	0.09386
Headtotal entry	13.075	40	0.68827	0.10883
Headtotal exit	14.0375	40	0.77944	0.12324

Materials and Methods

The study encompasses a total of n=40 children in primary school-aged 7-8, who were subjected to logopedic therapy aimed to overcome difficulties in graphomotor skills. It was conducted for a period of 5 years (2012-2017). The implemented therapy was done either individually or in groups of 2 children.

The children, part of this study who were subjected to logopedic therapy, after initial evaluation by a medical board were diagnosed with the following disorders according to ICD-10: F80.9 “Developmental disorder of speech and language, unspecified”, F81 “Specific reading disorder”, F81.3 “Mixed disorder of scholastic skills”, F81.8 “Other developmental disorders of scholastic skills. Developmental expressive writing disorder”, F81.9 “Developmental disorder of scholastic skills, unspecified” [5].

The number of children in this study was n=40, divided into males M-19 (47,5 %) and females F-21 (52,5 %) (Table 1).

Children from all age groups were evaluated before the initiation of the logopedic therapy with the following tests:

“Head test” (special postural praxis): Aimed for testing ideomotor praxis for new movements. In our study we used a version of the test with 13 movements.

“Test for evaluating written language in children with language deficits in primary school age”: The test consists of dictation. The texts used in this experiment were consistent with the age group of the children as well as the National school standards for the particular age group.

Data from the dictation were written in a protocol developed

Table 3:

Head test	Mean	Standard deviation	95% Confidence interval		t	Total	Significance
			Lowest	Highest			
En1- Ex1	-0.38750	0.40012	-0.51546	-0.25954	-6.125	40	0
En1.2- Ex1.2	-0.57500	0.52563	-0.74310	-0.40690	-6.919	40	0
EnTotal- ExTotal	-6.51250	0.59364	-6.70235	-6.32265	-9.384	40	0

Table 4:

Types of mistakes	Errors	Phonetic replacements	Graphic replacements	Additions	Repetitions	Rearrangements	Merging / Splitting	Others
Phoneme level	3,2%	54,5%	26,8%	6%	-	-	-	9,5%
Morphem level	3%	29,8%	20,2%	5,8%	3,8%	15,9%	11,5%	10%
Lexeme level	2,5%	3,7%	14,7%	4,2%	2%	22,7%	39,4%	10,8%
Syntaxeme level	1%	3,9%	13,6%	3,5%	5,3%	22%	40,6%	10,1%

by Yakimova (2009) [6] and for a certain number of mistakes an additional evaluation scale was added. The types of mistakes noted in the protocol were grouped in three levels: grapheme, morpheme as well as optically, phonetic-phonematic mistakes and misspellings. The work with the students continued for one year and in the end of the period they are being evaluated again with the described test for the particular age group.

For summarizing the results before initiating logopedic therapy and in the end of the school year the following statistical tools for analysis were used:

Descriptive statistic- Measurements of central trend, row width, error rate.

Correlation analysis- Relation of data.

T-test- Mean value comparison in different conditions of the dependent variable.

Results

Descriptive statistics of the compared methods in the studied group: Test sample number 1 “Head Test” (Table 2).

Identification of correlation between entry and exit “Head- test” results.

The statistical analysis of the compared methods in the 7-8 years age group confirms the described results and shows a significant increase of the total result of both parts of the test after implementation of logopedic therapy.

Significant increase of correctly done tasks from both sides of the test in our study can be explained by the neuropsychological basis of the processes related to increasing the efficacy of selective visual attention according to “response time” index typical for age 7-8 year olds.

The results gathered after implementation of T-test for comparison of the mean value in different conditions of the dependent variable for the indicators in “Head test”, represented in Table 3 after comparison, show significant difference in the two parts of the test study.

Those differences can be contributed to the fact that in the second part of the test (upon instruction) the children subjected to our study, 7-8 year olds have better level of formation of graphomotor skills.

Test sample 2 - Dictation

Table 4 represents percentage distribution of the types of mistakes on the different levels according to incidence rate at entry level.

Table 5:

Types of mistakes	Errors	Phonetic replacements	Graphic replacements	Additions	Repetitions	Rearrangements	Merging / Splitting	Others
Phoneme level	3,2%	54,5%	26,8%	6%	-	-	-	9,5%
Morpheme level	3%	29,8%	20,2%	5,8%	3,8%	15,9%	11,5%	10%
Lexeme level	2,5%	3,7%	14,7%	4,2%	2%	22,7%	39,4%	10,8%
Syntaxeme level	1%	3,9%	13,6%	3,5%	5,3%	22%	40,6%	10,1%

Table 6:

Methods	Mean	Total	Standard deviation	Standard error of the mean
Test 2 entry	14.9750	40	3.60546	0.57007
Test 2 exit	19.5000	40	3.36650	0.53229

Table 7:

	Total	Correlation	Significance
Test 2 entry & Test 2 exit	40	0.390	0.013

Table 8:

Test 2	Mean	Standard deviation	95% Confidence interval		t	Total	Significance
			Lowest	Highest			
EnT2 - ExT2	-4.52500	3.85631	-5.75831	-3.29169	-7.421	40	0

The results can be interpreted from the psychophysiological and neuropsychological model of initiation of the act of writing.

Underdevelopment of the complex manifestation of the visual-motor coordination in children with language deficits leads to difficulties in the already mentioned linguistic level.

Column "Others" encompasses a sum of punctuation mistakes and misspellings, poor level of literacy, lack of knowledge on graphemes. The percentage is relatively uniform in all linguistic levels.

Table 5 represents the percentage distribution of the types of mistakes on the different levels according to incidence rate at exit level after one year of logopedic therapy.

The results at exit level of the test study show positive influence of the therapeutic methods that we used. The difference of almost 20% drop of the "rearrangements and merging/splitting on lexeme and syntaxeme level" shows that the age of 8 years in children with language deficits is crucial for the formation of normal language functions. It is important to point out also the drop of nearly 20% of phonetic and graphic replacements on phoneme, morpheme and lexeme levels.

Descriptive statistics of compared methods in Test 2 (Table 6).

Identification of relation between entry and exit levels (Table 7).

T-test for comparison of mean values in different conditions of the dependent variable according to the indicators in Test 2 (Table 8).

Also in the results of Test 2, showed in (Table 6-8) significant correlations were marked at entry and exit levels and in the three indicators- compared methods, relations between entry and exit levels, comparison of the mean values indifferent conditions of the dependent variable of the indicators.

Discussion

Results show that the group of 7-8 years old children with language deficits is the sensitive moment which marks the start of formation with higher rate and maturation of structures of the

left hemisphere: frontoparietal and basal temporal, responsible for development of spatial orientation and the ideomotor praxis. Hence the implementation of intensive logopedic therapy aimed for strengthening and development of the affected components related to visual- motor organization is crucial for developing graphomotor skills and establishing and development of the written form of language.

Similar data has been published by Machinskaya, Krupskaya (2011) [7]. Other authors who study the visual gnosis within children also describe the age of 7-8 years as crucial for maturation of the executive control and increasing of the activity of structures, responsible for the ventral visual system [8-12]. Our results also confirm the increased active attention and maturation of the visual gnosis within children in that particular age group.

The origin and existence of phonetic replacements in certain linguistic level can be attributed to deficits of different origin of auditory perception and articulation disorders. The presence of high percentage of phonetic replacements in dictation as well as splitting in writing, seen within words constituted by a lot of consonants strongly inappropriate and hard to be articulately combined, are described by other authors [13,14].

Clinical studies of Bulgarian children with language deficits show that they have difficulty with: 1) perceptive tasks for differentiating phonemes in accordance to given feature (voiced/voiceless, sibilant/rustling sonority); 2) tasks for phonetic analysis as segmentation of phonemic numbers in words (quantitative sound analysis); defining the place of a given sound in words (quality sound analysis); separation of the first/last sound of words; discriminating different phonemes in paired words, that differ only by one word [15,16]. Phonological deficit is often associated with deficits in verbal memory and sometimes with difficulty in processing of purely verbal information according to instruction.

According to the data in English literature serious deficit in the area of morphology is observed in children with specific language disorders [17-20]. Even though they manage to master the grammatical morphemes in consistency similar to that of normal children, language deficit children show atypical types of control on those morphemes in time of production [15].

We believe that implementation of intensive logopedic therapy aimed for establishment and development of the deficient components related to visual-motor organization in this particular age is very important for development of graphomotor skills and formation and development of the written form of language.

Conclusion

The results of the implemented test samples in the age group of 7-8 years show that in children with language deficits this age period is sensitive for developing structures in left hemisphere and its frontal lobes, which are responsible for development of visual-motor

coordination but also for establishment of verbal praxis, speech formation and perception.

The implemented therapeutic techniques in the period of logopedic therapy are aimed not only for establishing visual- special orientation but also for formation and development of language competence which is related to developing the written form of language.

References

1. Raichev R, Raicheva M, Rasheva M, Matanova M, Stoyanova K, Raichev I. *Neuropsychology. Basis of neuropsychology*. 2012.
2. Mavlov L. *Fundamental Neurology*. 2000.
3. Peicheva S. *Cognitive Neuropsychology*. 2000.
4. Roux FE, Draper L, Kopke B, Demonet JF. Who Actually Read Exner? Returning to the Source of the Frontal "Writing Centre" Hypothesis. *Cortex*. 2010; 46: 1204-1210.
5. World Health Organization. *International Statistical Classification of Diseases and Related Health Problems. Tenth Revision*. 2003.
6. Yakimova P. *Disorders in written language Part 1*. 2009.
7. Machinskaya RI, Krupskaya EV. Ripening Regulatory Structures Of The Brain And Organizing Attention In Children Of Younger School Age. 2011; 4: 32-43.
8. Cycowicz YM. Memory development and event-related brain potentials in children. *Biological Psychology*. 2000; 5: 145-174.
9. Dukette D, Stiles J. The effects of stimulus density on children's analysis of hierarchical patterns. *Developmental Science*. 2001; 4: 233-251.
10. Farber DA, Petrenko NE. Individual Features Of Visual Recognition In Preschool Children. 2012; 1: 31-34.
11. Jolly C, Gentaz E. Analysis of cursive letters, syllables, and words handwriting in a French second-grade child with developmental coordination disorder and comparison with typically developing children. *Front Psychol*. 2013; 4: 1022.
12. Oliveira RE, Billington J, Wann JP. Optimal use of visual information in adolescents with developmental coordination disorder. *Exp Brain Res*. 2014; 232: 2989-2995.
13. Jekov J. Predictors of early diagnosis of dysgraphia in preschool and primary school age. *Dissertation*. 2010.
14. Richards TL, Grabowski P, Boord TJ, Yagle K, Askren M, Mestre Z, et al. Contrasting brain patterns of writing-related DTI parameters, fMRI connectivity, and DTI-fMRI connectivity correlations in children with and without dysgraphia or dyslexia. *Neuroimage Clin*. 2015; 8: 408-421.
15. Todorova E. *Dyslexia*. 2007.
16. Shtereva K. Structure of phonologic maturation in Bulgarian children. *Psychological Studies*. 2012; 1: 25-36.
17. Sela I, Karni A. Differences in learning volitional (manual) and nonvolitional (posture) aspects of a complex motor skill in young adult dyslexic and skilled readers. *PLoS One*. 2012; 7: e43488.
18. Sturm JM, Rankin-Erikson JL. Effects of hand-drawn and computer-generated concept mapping on the expository writing of middle school students with learning disabilities. *Learning Disabilities Practice*. 2002; 17: 124-139.
19. Stoeckel RE, Colligan RC, Barbaresi WJ, Weaver AL, Killian JM, Katusic SK. Early speech-language impairment and risk for written language disorder: A population-based study. *J Developmental Behavioral Pediatrics*. 2013; 34: 38-44.
20. Visscher C, Houwen S, Scherder EJA, Moolenaar B, Hartman E. Motor profile of children with developmental speech and language disorders. *Pediatrics*. 2007; 120: 158-163.